

PROGRESS OF FORENSIC ENTOMOLOGY RESEARCH IN INDIA FROM PAST TO PRESENT

Rita Kumari

(Research Scholar)

Department of Zoology

LNM University Darbhanga

ABSTRACT

Research on Forensic Entomology was started in 1889 in India. Several workers in India experimented on this line but the field has much more to develop. The aim of the present review is to describe different dimensions of research like Carrion Ecology Study, Forensic Entomotoxicology Study, Aquatic Forensic Entomology Study, Forensic Entomology research using morphological approach, Forensic Entomology research using molecular approach, forensically important flies on weather factors as well as seasonal variation.

Keywords: Forensic entomology research; India; review.

1. INTRODUCTION

Forensic entomology is the analysis of insect evidence for forensic and legal purposes [1]. Forensic entomology was first depicted in the thirteenth century in China by a lawyer cum death investigator Sung Tz'u in medico-legal textbook 'His yuan chi lu' (possible translation 'the washing away of wrongs'). Later on, Hoffman [2] reported insects and other arthropods as forensic indicator during mass exhumations in Germany and France. J. A. Payne [3] was supposed to be the first field entomologist of forensic entomology who performed an experiment on Pig (*Sus scrofa* L.) carrion during the summers of 1962 and 1963 in South Carolina and observed 522 arthropod species belonging to four Orders of Coleoptera, Diptera, Hymenoptera, and Araneida. Few experiments on this line were carried out by some enthusiastic persons in India. Dr. S. Coull Mackenzie, a Police Surgeon of the then Calcutta, was supposed to be the pioneering work in the field of forensic entomology in India. He described eight cases of saponification of human dead bodies in and around Calcutta in 1889. No further work on this aspect was reported after Mackenzie's report [4]. With the establishment of Central Forensic Science Laboratory at Kolkata in 1957, research on Forensic Entomology was developed slowly. Several workers from that time have performed different kind of researches under the broad field of Forensic Entomology up to recent times that may be categorised into six different dimensions like:

- a) Carrion Ecology Study,
- b) Forensic Entomotoxicology Study,

- c) Aquatic Forensic Entomology Study,
- d) Forensic Entomology research using morphological approach,
- e) Forensic Entomology research using molecular approach,
- f) Forensically important flies on weather factors as well as seasonal variation.

All the segments pointed out in the present review work, fall under the vast area of research of forensic sciences related to entomology. Undoubtedly every segment of study aims to investigate postmortem interval very minutely following its methodology which is described in brief to illuminate the affair.

a) Carrion Ecology Study: It is the entomofauna study on a decomposing animal or human dead body. In the field of Carrion Ecology, larval stages, pupal stages of different developmental maturity and adults from human dead bodies or animal carcasses are usually collected. Various stages of decomposition viz. fresh, bloat, active decay, advanced decay and dry remains of the corpse are observed. Some workers mentioned a detailed list of insects involved in each phase of decomposition. Developmental stages of maggot were observed by Rao et al. [5] to measure the time since death. With increasing interest of the study of insect behaviour on cadavers, several researchers like Kulshrestha and Chandra [6] studied various environmental conditions and faunal role on a dead body. Antemortem wounds which became unidentifiable were also observed by them. The authors described 25 cases of cadavers related to maturation of blowflies. In another investigation, Kulshrestha and Satpathy [7] evaluated Post Mortem Interval (PMI) by collecting beetles *Dermestes maculatus* (Coleoptera, Dermestidae) and *Necrobia rufipes* (Coleoptera, Cleridae) from two dry female skeletal remains. In a sensational death case of Exminister of Karnataka, probable PMI was determined by Kulshrestha and Satpathy [8] examining samples of fly eggs, first instar and second instar maggots on a dead body. The entomological method was superior and statistically more reliable than other medico-legal approaches in the estimation of PMI that was proved by Kashyap and Pillay [9] by examining sixteen insect-infested cadavers. Various cases of insect succession were studied by many workers. Different species of blowflies (Diptera, Calliphoridae) viz. *Calliphora vicina*, *C. vomitoria*, *Lucilia illustris*, *Chrysomya megacephala*, *C. rufifacies* were collected by Singh and Bharti [10] from rabbit carcasses in the field and various putrefying cadavers in many places of Punjab in different seasons. In another study, eight species of ants viz. *Camponotus compressus*, *Crematogaster contemta*, *C. hodgsoni*, *Dorylus labiatus*, *Meranoplus bicolor*, *Messor barbarus*, *Paratrechina longicornis*, *Tapinoma melanocephalum* (Hymenoptera, Formicidae) were observed by Singh and Bharti [11] from rabbit carrion in the field during various seasons of the year.

Three species of Muscidae and one species of Sarcophagidae were recorded by Bharti et al. [12] from India for the first time from carrion of rabbit in Patiala campus of the Punjabi University of state Punjab. Muscids viz. *Hydrotaea occulta*, *H. capensis*, *Atherigona savia* and *Sarcophaga princeps* were recorded. To show nocturnal oviposition behaviour of blowfly, an experiment was conducted by Singh and Bharti [13] around fourteen night trials at an unilluminated area in Patiala by taking mutton as a bait on the top of a pole and 3 types of Calliphorids (*Calliphora vicina*, *C. megacephala*, *C. rufifacies*) laid eggs in five cases. It was rather an extension of an experiment designed by Greenberg [14]. A similar type of study was performed by Singh and Bharti [15] at an unlighted area in Patiala, Punjab taking a piece of mutton as bait on the top of a pole to observe nocturnal larviposition behaviour of flesh flies in September 2005. It was found that *Sarcophaga albiceps* and *S. hirtipes* were able to deposit their larvae at night also. In an elaborated successional study, a total of 38 insect species belonging to four orders and thirteen families (Calliphoridae, Sarcophagidae, Muscidae, Anthomyiidae, Otitidae, Staphylinidae, Histeridae, Cleridae, Dermestidae, Tenebrionidae, Silphidae, Formicidae, Tineidae) were collected by Bharti and Singh [16] on decaying rabbit carcasses and it was surveyed that maximum diversity of insect species occurred in Spring. Twenty cases of death were studied by Aggarwal et al. [17] to monitor the insects involved in the decay of corpses. Gupta and Setia [18] suggested that Forensic entomology was utilised to determine PMI, the season of death, geographical location of death, movement or storage of remains after death etc. and molecular identification was also exposed a new arena in that field. Singh and Sharma [19] criticised forensic research in India. They remarked that lack of baseline data and experienced professionals for measurement of minimum post-mortem interval made the line most untouched till date. The baseline data were mainly involved in taxonomy, zoogeographic distribution, biology and ecology of insects. Singh and Bala [20] surveyed the time length of the starvation period required to induce post feeding behaviour shortened with increasing age of larvae.

A brief overview of the necessity of forensic entomological data in a criminal investigation was provided by Joseph et al. [21]. These data were related to insects and their larval morphology, growth histories, species distribution and toxic contents in their tissue. An initiative was taken by Goyal [22] to measure the postmortem interval observing members of Calliphoridae, Sarcophagidae, Muscidae, Silphidae, Dermestidae at different decomposition stages on 47 human corpses in Amritsar, Punjab. The PMI of a headless 9 months old male foetus was calculated by Babu et al. [23] finding in the forest nursery at Jagdalpur on the basis of the developmental period of second instar larvae of blowfly *C. rufifacies* utilising Accumulated Degree Hours(ADH) method. The death time duration of human corpses was estimated by Chandna [24] measuring length of maggots of house fly (*Musca domestica*), flesh fly (*Sarcophaga* sp), skipper fly (*Piophilidae*) (*Piophilidae*) and blowfly (*Calliphora vomitoria*) with scale in millimetres in Karnal, Haryana. Bala and Kaur [25] observed insects from buried remains. For that purpose, an experiment was conducted on a buried piece of pork at a depth of 30 cm around 16 days in Ghawaddi village of Ludhiana in the state of Punjab and ten species of beetles belonging to 6 families and 2 species of order Hymenoptera were observed by them. Insect succession on the same piece of pork was observed by Bala and Kaur

[26] that was buried in the soil after 48 hrs of insect exposure in the same place. In that case, 41 insect specimens belonging to 13 species under 9 families of the order Diptera, Coleoptera and Hymenoptera were monitored. Singh et al. [27] criticised that many research papers and review articles remained about terrestrial forensic entomology, but few numbers were available on buried dead bodies. So, they reviewed the works to estimate post-burial interval, its importance and future perspective. A case study was done by Bala and Sharma [28] on a decapitated 38 years aged male human dead body at Punjab. Few third instar larvae of *C. megacephala* were collected from the amputated arm of the corpse, and they calculated the number of days required for *C. megacephala* to develop from egg to larvae. Tentative PMI was estimated at 10.6°C and according to them, that was 9 days. Bharti [29] had taken an initiative to find out the correlation between faunal composition and change in altitude in Himalayan regions. For that purpose, a study was conducted at a different elevation in North West Himalaya using carcasses of a cow as bait.

It was reported that blowfly diversity was decreased rapidly at an altitude of 2511m showing only three species due to limited abiotic factors and *C. villeneuvi* was abundant at that elevation. *C. megacephala* and *C. rufifacies* were monitored at first three elevations (350m, 970m and 2057m) because having extreme tolerance capacity of temperature fluctuations and flies belonging to genus *Calliphora* were present at higher altitudes of 2057m and 2511m because those were cold resistant. In a similar type of study, 19 species of blowflies belonging to *Chrysomya*, *Lucilia* and *Calliphora* were observed by Jadav and Sathe [30] in the Western Ghats. They also reported that the number of *Lucilia* and *Chrysomya* were more in low altitude. The authors further analyzed that diversity of blowflies was decreased with increasing altitude. A study was performed by Singh et al. [31] on thirtyeight human corpses to estimate the PMI using statistical tool SPSS version 20. It was observed that time duration for egg laying was 1-3 days. In another study, 25 species of forensically important 12 families of order Diptera were reported by Sathe et al. [32] from Western Maharashtra, India using pig liver as bait or from dead bodies. Later, the succession and life cycle of *Dermestes maculatus* was studied on exposed piglets' carcasses by Sonker et al. [33]. Fly specimen was abundant during spring and summer. In a different study, *S. (Liosarcophaga) dux* was reported by Chakraborty et al. [34] for the first time in India on a corpse of house gecko *Hemidactylus flaviviridis*. Another experiment was performed by Chakraborty et al. [35] on the cadavers of *Gallus gallus* around three different seasons including premonsoon, monsoon and post-monsoon to observe the effect of seasonal variation on development of immature stages of *S. (Parasarcophaga) albiceps* using one way ANOVA. It was found that there were no significant differences in the average length, average width and average biomass of the immature stages in seasonal variation. In the same year, another study was performed by Das et al. [36] on the carcasses of white rat (*Rattus rattus*) in West Bengal. They reported *Musca domestica*, *Lucilia cuprina* and *Hemipyrellia ligurriens* in it. Later, a study was conducted by Singh et al. [37] on laboratory-bred rat (*Rattus norvegicus*). Nine species of flies belonging to four families, viz. *Calliphoridae* (44.24%), *Sarcophagidae* (16.53%), *Muscidae* (28.87%) and *Phoridae* (13.35%) were observed and *P. ruficornis* firstly arrived and *M. domestica*, *C. megacephala* remain throughout the experiment, whereas *Megaselia* sp was present on the third day.

The PMI from larvae of *C. megacephala* was determined by Sharma et al. [38] using base temperature or lower threshold temperature in ADH method. In another work, Sharma et al. [39] suggested various methods for determination of PMI of Calliphorid flies and also analysed problems facing by forensic entomologists while measuring the time since death. Sharma and Singh [40] critically evaluated the progress of forensic entomology from past to present days emphasising PMI and insect succession in different countries adding Indian scenario also. Past and present overviews of India in the field of forensic entomology were given by Sharma and Singh [41]. The PMI of a mummified body of 23 year old female by observing the developmental time of *C. megacephala* from egg to pupa utilizing Accumulated Degree Hours (ADH) at $25\pm 1^{\circ}\text{C}$, was estimated by Bala and Sharma [42]. Their estimated PMI value was 9.6 days. A case study was done by Sharma and Bala [43] on 45 years aged male dead body in Ludhiana, Punjab, India. PMI was determined to utilise ADH method. Dynamics on human corpses by local insects like *C. megacephala*, *C. rufifacies*, *Dermestes maculatus*, *Necrobia rufipes*, *Phormia regina*, *C. albiceps* were observed by Sharma et al. [44].

b) Forensic Entomotoxicology Study: This is generally carried out when necessary tissues of corpses were not obtained to conduct toxicological study or at that situation when blood, urine or internal organs were absent for criminal investigation. A study on forensic entomotoxicology was conducted by Gola and Lukose to determine the presence of chemicals or drugs in insect body. For that purpose Librium (chlordiazepoxide), Diazepam, Prednisolone, Phenobarbitone and DDT were used as ante-mortem administered drugs or chemicals. They found the presence of Librium (chlordiazepoxide), Diazepam, Prednisolone and Phenobarbitone in all the three larval stages, pupal stages and adult form of blowflies but DDT was absent beyond the larval stages. Verma investigated life cycle pattern of *C. rufifacies* under four different temperatures like Cool temperature (Humid $20-24^{\circ}\text{C}$), Cool temperature (Dry $18-22^{\circ}\text{C}$), Room temperature (Humid $26-30^{\circ}\text{C}$) and Room temperature (Dry $24-28^{\circ}\text{C}$) and effect of two drugs ethanol and cannabis on the development of it. It was observed that higher temperature and humidity accelerated the growth and maturation. According to them, ethanol and cannabis showed faster development than the control condition. Singh et al. conducted an experiment by rearing *C. megacephala* on rat tissues administering half lethal (37.5 mg/kg bw), lethal (75 mg/kg bw) and twice lethal (150 mg/kg bw) dosages of Ketamine hydrochloride previously to monitor the effect on the development of fly taking different parameters like weight, length, width, time of adult emergence and survival rate. In another study, Singh and Bhupinderjit reared larvae of *C. megacephala* on rat tissues exposed to Cadmium chloride (CdCl_2) in three different concentrations viz. half lethal (3.25mg/kg bw), lethal (6.5 mg/ kg bw) and twice lethal (13 mg/kg bw) by intraperitoneal injection to observe the effect of Cadmium chloride on development of *C. megacephala*.

c) Aquatic Forensic Entomology Study: This is related to the estimation of PMI in case of carcasses found in the submerged condition in aquatic sources. Very few works on aquatic forensic entomology have done in India till recent time. Many papers were available on forensic entomology in terrestrial habitat but very little works were done in an aquatic habitat. A review paper about aquatic forensic entomology pointed out only its importance, future prospective and forensically important insects in the aquatic environment were published by

Sharma and Singh. Another experiment on this line was designed by Singh and Greenberg by submerging pupae of known ages of five species of blowflies one to five days duration underwater to know chances of survival. It was observed that 25% of pupae of *Protophormia terraenovae*, *Phormia regina*, and *Phaenicia sericata* (Diptera, Calliphoridae) generated normal adult flies after 4 days of immersion but none after 5 days. They also reported the expansion of pupal period among survivors occurred in relation with duration of submersion. According to Singh et al. maggots of *C. megacephala*, Sarcophagids and diatoms conjointly helped in PMI estimation.

d) Forensic Entomology research using morphological approach: Accurate morphological identification of immature or adult stages of insects indicates one of the paramount factors for forensic investigation because it indirectly helps us to determine postmortem interval. Identification of species of Calliphoridae was done by Greenberg and Singh through analysing the morphology of eggs of its 11 species belonging to six genera with SE study. Ultrastructural study of first, second, third instars and puparium of *P. ruficornis* were observed by Singh et al. using light microscopy and scanning electron microscopy for the first time in India. Main diagnostic characters, i.e. cephalopharyngeal apparatus, cephalic segment, structure and orientation of spines, pupal respiratory horns, structures of both anterior and posterior spiracles were considered by them. Those characters are certainly helpful in forensic investigations of fly species. Bala and Sharma commented that traditional techniques of PMI of blowflies in most cases were accurate but lack of precision. So, they mentioned some recent techniques of blowfly age estimation that was pteridine fluorescence analysis, cuticular banding pattern analysis, internal morphological analysis, cuticular hydrocarbon analysis, gene expression analysis, volatine organic compounds analysis released by larvae and pupae. Sharma and Singh suggested various morphological and molecular bases of identification methods for forensically important Indian flies. According to them, morphological methods were a study of Allozymes, Cuticular hydrocarbon techniques and Scanning Electron Microscopy whereas techniques using Nuclear DNA, Nuclear Ribosomal DNA and Mitochondrial DNA as markers were included in molecular-based identification methods. In another case, SEM studies on the morphology of sensilla of antenna of adult male and female of *H. ligurriens* were observed by Hore et al. for the first time. That finding helped easier identification of insect morphology from the forensic entomological viewpoint and better distinction of *ligurriens* from other calliphorids.

e) Forensic Entomology research using molecular approach: Identification of insect specimens through morphological approach is confusing in some instances. Having similar characteristic features of some individuals, a parallel molecular-based identification method was evolved to simplify the identification process using various molecular markers like Cytochrome Oxidase subunit I (COI) and subunit II (COII), NAD Dehydrogenase subunit 5 (ND5), 16S rRNA. In India, molecular Study in the field of forensic entomology was done on Sarcophagid, Calliphorid and very little on Muscid flies using mitochondrial genes and 16S rRNA as a marker. According to Bajpai and Tewari two mitochondrial genes Cytochrome Oxidase subunit I (COI) and NAD Dehydrogenase subunit 5 (ND5) were helped to know genetic relatedness and to construct phylogeny among five sarcophagid species using PCR. In another

study, five sympatric species of the genus *Sarcophaga* was described to determine genetic relationship using mitochondrial COI and II region . Identification of forensically important fly *C. megacephala* and *C. rufifacies* was done to observe polytene chromosomes from trichogen cells remain in the pupae . Molecular identification of three Sarcophagid flies was carried out by Sharma et al. based on 450 bp region of Cytochrome Oxidase I gene. DNA amplification study was done by Khullar and Singh from 13 to 14 years old insect samples of four species viz. *C. vomitoria*, *H. pulchra*, *Catapicephela splendens* and *Catapicephela ingens* (Diptera, Calliphoridae) using 16S rRNA primer. It was observed that COI gene was highly fragmented but the 16S rRNA gene remained well protected from fragmentation and gave ideal results for old samples.

f) Forensically important flies on weather factors as well as a seasonal variation: Temperature and Humidity have great roles in the development of insects. Season also shows pronounced impact on life cycle duration of insects. A study was done by Karunamoorthy and Lalitha in which they reared larvae of forensically important Australian sheep blowfly *Lucilia cuprina* on lean beef and adults on a glucose-nutrient broth mixture, lean beef and sugar. Life spans of both adult male and female were from 23 to 31 days at $32 \pm 2^\circ\text{C}$. Another study was conducted by Bharti and Singh on the carcasses of domestic rabbits to observe various larval stages of three blowfly species viz. *C. vicina*, *C. megacephala* and *C. rufifacies*. According to the authors, *C. megacephala* and *C. rufifacies* were found to breed around the year, whereas *C. vicina* was able to breed on carcasses only during winter and spring season. Then, the development time of *C. megacephala* at four different temperatures 22°C , 25°C , 28°C , 30°C at laboratory environment was investigated by Bharti et al. and it was reported that there was an inverse relationship between temperature and time of development. The time periods required to complete lifecycle of the fly were 15.5 days, 12.4 days, 8.5 days and 6.3 days at 22°C , 25°C , 28°C and 30°C temperatures respectively. Bharti [88] found a correlation between temperature and time required to complete life cycle of *C. vicina* and *M. domestica* and it was concluded that when the temperature was increased, days required to complete life cycle became shortened. The life cycle of *P. ruficornis* in the rainy season was studied by Patil. It was observed that maggots were required four days to reach pupal stage and pupa was converted to adult in eleven days. So, the whole life cycle of the fly was completed in sixteen days according to the author. It was also suggested that the life cycle study of *P. ruficornis* for PMI estimation was done scarcely in India than Europe, America and Australia. In another study, *C. megacephala* and *C. rufifacies* were

surveyed at four different temperatures (22°C , 25°C , 29°C and 31°C) in order to reach an interrelationship between larval age, body length and body dry weight.

2. DISCUSSION

Different dimensions of forensic entomological research are highlighted and from this discussion, it is evident that major forensic entomologists in India engaged themselves in Carrion Ecology study. Some workers focused more on the determination of post submergence interval and post-burial interval also. Pankaj Kulshrestha, the forensic specialist of

MedicoLegal Institute of Bhopal, is a pioneer in this area and various workers of Punjabi University also have done a tremendous job. Many workers also engaged themselves to observe the development of insects in consequence of temperature and humidity factors in association with seasonal variations. But other dimensions of Forensic Entomology like Forensic Entomotoxicology Study, Aquatic Forensic Entomology Study, Post-burial interval Study and forensic entomology research using morphological approach are not properly excavated. Forensic entomology research using molecular approach has been initiated in the recent decade and major portion of works has been done on Sarcophagids, Calliphorids and very little on Muscids. More emphasis should be given to forensic entomology research in India in future.

3. CONCLUSION

Forensic entomology study has many dimensions which were studied in India by different workers at different times. Research on this field has not been done up to the mark till date. Forensic entomology research using morphological approach has to be developed more in India.

ACKNOWLEDGEMENTS

The authors acknowledge the Teacher-in-Charge, Indas Mahavidyalaya for his encouragement and the Principal of Sonamukhi College for necessary permission and help in research work.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Amendt J, Campobasso CP, Gaudry E, Reiter C, LeBlanc HN, Hall MJR. Best practice in forensic entomology—standards and guidelines. *Int J Legal Med.* 2007;121:90–104.
2. Hofmann, O. Observations de larves de Dipteres sur des cadavres exhumes (Observations on Diptera larvae on exhumated corpses). *C.R. Séances Soc. Ent. Belg.* 1886; 74:131-132. (In French)
3. Payne JA. A summer carrion study of the baby pig *Sus scrofa* Linnaeus. *Ecol.* 1965;46:592-602.
4. Mackenzie SC. Eight cases of saponification. *Indian Med Gazette.* 1889;1889:42-44.
5. Rao MB, Goyal MK, Kushwaha KPS, Gaur JR. Experimental study of the developmental stages of maggots for estimation of time since death. *J Forensic Sci.* 1984;24(4):398.
6. Kulshrestha P, Chandra H. Time since death. An entomological study on corpses. *Am J Forensic Med Pathol.* 1987;8(3):233-8.

7. Kulshrestha PK, Satpathy DK. Use of beetles in forensic entomology. *Forensic Sci Int.* 2001; 120(2001):15-17
8. Kulshrestha PK, Satpathy DK. Forensic entomology analysis professional help in Karnataka case. *JIAFM.* 2005;27(2):117-119.
9. Kashyap VK, Pillay VV. Efficacy of entomological method in estimation of postmortem interval: A comparative analysis. *Forensic Sci Int.* 1989;40(1989):245-250.
10. Singh D, Bharti M. Forensically important blowflies (Diptera: Calliphoridae) of Punjab (India). *Uttar Pradesh J Zool.* 2000;20(3):249-251.
11. Singh D, Bharti M. Ants (Hymenoptera : Formicidae) associated with decaying Rabbit carcasses. *Uttar Pradesh J Zool.* 2001;21(1): 93–94.
12. Bharti M, Singh D, Sidhu IS. First record of some carrion flies (Diptera : Cyclorrhapha) from India. *Uttar Pradesh J Zool.* 2001;21(3): 267-268.
13. Singh D, Bharti M. Further observations on the nocturnal oviposition behaviour of blow flies (Diptera: Calliphoridae). *Forensic Sci Int.* 2001;120:124-126. DOI: 10.1016/S0379-0738(01)00419-4
14. Greenberg B. Nocturnal oviposition behaviour of blow flies (Diptera: Calliphoridae). *J Med Entomol.* 1990;27(1990):807-810.
15. Singh D, Bharti M. Some notes on the nocturnal larviposition by two species of Sarcophaga (Diptera : Sarcophagidae). *Forensic Sci Int.* 2008;177(2008):e19-e20.
16. Bharti M, Singh D. Insect faunal succession on decaying rabbit carcasses in Punjab, India. *J Forensic Sci.* 2003;48(5):1-11.
17. Aggarwal AD, Gorea RK, Aggarwal OP, Singh D. Forensic entomology - a guide to postmortem interval. *JPAFMAT.* 2003;3:7-10.
18. Gupta A, Setia P. Forensic entomology -past, present and future. Anil Aggrawal's Internet J Forensic Med Toxicol. 2004;5(1):50-53.
19. Singh J, Sharma BR. Forensic entomology: A supplement to forensic death investigation. *JPAFMAT.* 2008;8(1):26-33.
20. Singh D, Bala M. The effect of starvation on the larval behavior of two forensically important species of blow flies (Diptera : Calliphoridae). *Forensic Sci Int.* 2009; 193(2009):118-121.
21. Joseph I, Mathew DG, Sathyan P, Vargheese G. The use of insects in forensic investigations: An overview on the scope of forensic entomology. *J Forensic Dent Sci.* 2011;3(2): 89-91.

22. Goyal PK. An entomological study to determine the time since death in cases of decomposed bodies. *J Indian Acad Forensic Med.* 2012;34(1):10-12.
23. Babu BS, Sharma H, Bharti M. Estimation of post mortem interval by rearing *Chrysomya rufifacies* (Macquart, 1842) (Diptera: Calliphoridae): A case study from central India. *Anil Aggrawal's Int J Forensic Med.* 2012; 14(2):12. Available: http://anilaggrawal.com/ij/vol_014_no_002/papers/paper002.html
24. Chandna SS. Insects in forensic science for detection of crime. *New York Sci J.* 2012; 5(2):27-31.
25. Bala M, Kaur P. Insect faunal succession on buried piece of pork in the State of Punjab (India): A preliminary study. *J Forensic Res.* 2014;6(1):1-4. DOI: 10.4172/2157-7145.1000252
26. Bala M, Kaur P. Entomofauna on decomposed piece of pork: Study on delayed burial. *J Ent Res.* 2015;39(1):77-84.
27. Singh R, Sharma S, Sharma A. Determination of post burial interval using entomology: A review. *J Forensic Legal Med.* 2016;42(2016): 37-40.
28. Bala M, Sharma A. A case study of decapitated 38 years old male corpse from Punjab (India): Image Article. *J Forensic Res.* 2015; 6(2). DOI: 10.4172/21577145.1000i102
29. Bharti M. Altitudinal diversity of forensically important blowflies collected from decaying carcasses in Himalaya. *Open Forensic Sci J.* 2012;5:1-3.
30. Jadav D, Sathe TV. Altitudinal diversity of forensic blow flies (Diptera: Calliphoridae) of Western Ghats (Maharashtra). *J Forensic Res.* 2014;5(6):1-4.
31. Singh H, Venketasan M, Aggarwal OP, Raj S. Use of maggots for the estimation of time since death. *J Punjab Acad Forensic Med Toxicol.* 2014;14(1):22-26.
32. Sathe TV, Sathe A, Sathe TN. Diversity of dipterous forensic insects from Western Maharashtra, India. *Int J Pharm Bio Sci.* 2013; 4(2):173-179.
33. Sonker R, Rawat S, Singh K. Succession and life cycle of beetles on the exposed carcass. *Int J Sci Innov Res.* 2015;3(1):46-50.
34. Chakraborty A, Ansar W, Ghosh S, Banerjee D. The first report of the life cycle of *Sarcophaga (L.) dux* on dead reptilian carcass: Their application as forensic indicators. *Sch Acad J Biosci.* 2014;2(11):731-739.
35. Chakraborty A, Ghosh S, Ansar W, Banerjee D. Developmental analysis of immature stages of *Sarcophaga (Parasarcophaga) albiceps* Meigen, 1826 (Diptera : Sarcophagidae) on *Gallus gallus* carcass: Their applications as forensic indicators. *IOSR J Agric Vet Sci.* 2015;8(8):79-89.
36. Das P, Majumder S, Imam I, Mallick K, Bhattacharya M. Additional record of some

forensically important Dipteran Species from West Bengal. *Bionotes*. 2015;17(2).

37. Singh N, Kumari V, Singh NP. An analysis of the ecological succession pattern of diptera on the carcass of laboratory bred rats. *Int J Agric Innov Res*. 2016;4(5):837-841.

38. Sharma R, Garg RK, Gaur JR. Estimation of post mortem interval from larvae of *Chrysomya megacephala* (Fabricus) using the concept of ADH. *Indian Police J*. 2013;200-212.

39. Sharma R, Garg RK, Gaur JR. Various methods for the estimation of the post mortem interval from Calliphoridae: A review. *Egyptian J Forensic Sci*. 2015;5:1-12.

40. Sharma M, Singh D. Historical approach of forensic entomology: A review. *Int J Adv Res Sci Eng Technol*. 2015;2(9):838-845.

41. Sharma M, Singh D. Forensic entomology: An Indian prospective. *Int J Sci Res Sci Eng Technol*. 2016;2(1):207-209.

42. Bala M, Sharma A. Postmortem interval estimation of mummified body using Accumulated Degree Hours (ADH) method: A case study from Punjab (India). *J Forensic Sci & Criminal Inves*. 2016;1(1):1-5.

43. Sharma A, Bala M. Case study and PMI estimation of male corpse from Ludhiana, Punjab, India: An implication of ADH method. *Indian J Forensic Med Toxicol*. 2016;10(2): 28-33.

44. Sharma A, Bala M, Singh N. Five case studies associated with forensically important entomofauna recovered from human corpses from Punjab, India. *J Forensic Sci & Criminal Inves*. 2018;7(5):1-6. DOI: 10.19080/JFSCI.2018.07.555721006