

**EFFECT OF EDOSULFAN ON THE BLOOD GLUCOSE LEVEL IN THE INDIAN
LIZARD CHAMELEON ZELNICUS.**

Dr. Bishnu Charan Pradhan

Dept. of zoology

Angul Mahila Mahavidyalaya, Angul Odisha. 759122

India

ABSTRACT

The present study is aimed to investigate the effect of Endosulfan on the blood of Indian lizard, Chameleon zelenicus. In the present study, the level of blood glucose was observed on the median lethal dose of endosulfan (42.7 mg/kg body weight) for 6,12,18,24,48,72,96 and 120 hours of exposure and lethal dose of (59.9 Endosufan m g/kg body weight) for 6,12 and 18 hours of exposure. The present study showed the level of blood glucose was decreased in the blood of indian lizard,. Chameleon zelenicus

Key words: Endosulfan , Blood Glucose, Chameleon zelenicus .

1.INTRODUCTION

A wide range of toxic chemicals called pesticides are used to control or eradicate the pests of agriculture, horticulture and other different crops throughout the world by people. The uses of pesticides have served to improve our crops from the ravages of insect pests. Control of pests have always been one of the major factors in enhancing the agricultural production (Hague and Freed¹⁹, 1975)

The synthetic group of pesticides are broadly-classified into three major groups viz., Organochlorines, Organophosphorus and Carbamates (Corbett,¹¹ 1973). Among them organochlorine pesticides are commonly used in agriculture due to their high effectiveness in reducing the pests even at low concentrations. The organophosphorus compounds dichlorvos, monocrotophos, parathion, fenitrothion, malathion, endosulfan, phosalone, dimethoate, thiometon are degradable and are quickly broken down into harmless products in the milieu. Though they are highly toxic, they are not pollutants if used carefully, whereas the carbamate compounds isolan, carbaryl, carbofuran, aldicarb, carbathion are least in their toxicity. Among the organochlorine compounds DDT, aldrin, lindane, dieldrin, endrin, chlordane and heptachlor are remarkably long lasting chemicals in the milieu. For instance, residues of DDT and some

other pesticides have been reported to be found even after 12 years from their day of application and thereby causing biomagnification.

Blood plays an important role in the normal functioning of the body of any organism. Seasonal variations in the blood contents of reptiles have been reported by Hutton and Goodnight¹¹ (1957), Binyon and Twigg⁵ (1965), Kekic²⁷(1970). Banerjee and Banerjee⁴(1969) and Choubey⁹ (1975)

have also reported the seasonal variations in the blood components of tropical reptiles. Jayaraman²⁵(1976) has studied on the modulation of humoral and cell mediated responses to sheep erythrocytes in the lizard, Chameleon zelenicus. The influence of post helminth infection on haematological indices of Chameleon zelenicus was reported by Kameswari and Narsimha Rao²⁶ (1979). Banerjee (1981) has reported the seasonal variations of some blood constituents of the Indian lizard Chameleon zelenicus.

Blood glucose levels have been reported to be a sensitive indicator of many stresses viz., handling, forced activity, salinity changes and few chemical pollutants (Ghosh¹⁸ et al., 1969; Wedemayer,⁵¹ 1972; Silbergeld,⁴⁵ 1974 and Watson and Mckeown,⁴⁹ 1976.) A definite seasonal variation in the blood glucose level has been reported to occur in reptiles (Dessauer,^{14,15} 1953). Pancreatectomy affects the blood glucose differently in different reptiles (Houssay and Penhos²¹, 1960; Madhavan and Rangnekar, 1985). The effect of Cad₂ on the cells of endocrine pancreas in the snake, *Natrix natrix* was worked out by Calugrenan⁶ (1970). Light and electron microscopic studies on pancreatic islets of the lizard, *Lygosoma laterale* was observed by Rhoten and William⁴³ (1971). Hussaini²² and Amani (1976) studied the effect of intramuscular injections of two different doses of alloxan on the pancreatic islet tissue and the blood sugar level of the toad, *Bufo regularis* and the lizard, *Agama stillio* at different time intervals.)

An interesting study was made by Chan⁷ et al. (1970) on the water and electrolyte composition of the Iguanid lizard, *Dipsosaurus dorsalis* with reference to the control in the pituitary gland and the adrenal cortex. Lofts²⁹ et al. (1971) have observed the seasonal changes in the histology of the adrenal gland of the Cobra, *Naja naja*. Unsicker⁴⁷ (1974) has investigated the innervation of adrenal cells in the lizards, *Lacerta dugesi* and *Lacerta ptyusensis*. Contiguity of the adrenaline -s-rorir.g chromaffin cells with the interrenal tissue in the adrenal gland of a lizard was worked out by Conte¹⁰ (1977).

2. MATERIALS AND METHODS

Procurement, acclimatization and selection of the experimental animals

The experimental animals were live trapped in and around Angul area. The collected experimental animals were lodged in cages of 18 x 18 x 45 cms size and were acclimated for a week. They were nourished with live cockroaches, grasshoppers and butterflies. Water was placed ad libitum. Only the female animals of Chameleon zelenicus were taken for the present study. Healthy individuals weighing about 30 - 37 gms were selected for the control and the experimental studies. Much care was taken to avoid injured, pregnant and infected animals.

Chemical and physical properties of the selected pesticide

Endosulfan is organophosphorous pesticide extensively used to control a wide range of sucking and chewing pests of field crops, fruits and vegetables. It has many structural similarities with naturally occurring compounds, and their primary target of action in insects is the nervous system; it also inhibits the release of the acetylcholinesterase at the synaptic junction (Cabello et al 2001). Acetylcholinesterase plays a key role in the control of nerve excitability at post synaptic sites. Endosulfan inhibits the acetylcholinesterase. Inhibition of liver acetyl cholinesterase (AChE) activity is generally regarded as a useful indicator of poisoning by. Before this, lethal toxicity tests were carried out for four different concentrations of the endosulfan.

Toxicity studies

The toxicity tests have been classified into different categories based on the types of pollution in the environment as preliminary screening tests and tests to establish water quality criteria, etc. In the laboratory, determinations of toxicity may be divided into two types viz., short term or acute toxicity tests and long term or chronic biological responses (Negilski,³⁹ 1975).

Range finding tests

To avoid much time delay and effort the preliminary exploratory tests were conducted to get a range but not an accurate dose of the pesticide, which should be covered in the full scale tests. To conduct these tests, 1% stock solution of the pesticide, endosulfan was prepared by dissolving it in the coconut oil. Test solutions of required concentrations were prepared from the stock solution.

Lizards were orally administered with 1.0 ml of the pesticide solution, containing the required dose of pesticide by using hypodermic syringe without the needle. Much care was taken to avoid

regurgitation of the pesticide. The exploratory tests showed the range of toxicity for this pesticide *ho* this experimental animal between 47.7 to 69.7 rag/kg body weight.

Full scale tests

Based on the exploratory tests, full scale tests were conducted to assess the acute toxicity of the selected pesticide,. In endosulfan these tests seven concentrations viz., 42.7, 44.7, 47.7, 50.7, 53.7, 56.7p.nd 59.9 mg/kg body weight were administered and for each concentration six lizards were used. Simultaneously the control animals were maintained and they were administered with 1.0 ml of coconut oil by using hypodermic syringe without the needle. The control and experimental lizards were observed for 120 hours. None of the control individuals attained mortality. In the pesticide treated lizards, the mortality rate of the lizards were observed at different hour's viz., 6, 12, 18, 24, 48, 72, 96 and 120. The dose at which 50% mortality attained for a period of 120 hours was taken as median lethal dose (120 hours, LD₅₀) and the dose at which cent percent mortality attained in the test animals for a period of 24 hours was taken as lethal dose (24 hours, LD₁₀₀) for the present study.

From the observed results, it was inferred that the pesticide concentration of 42.7 mg/kg body weight at which 50% mortality occurred for 120 hours was taken as median lethal value (120 hours, LD₅₀) and the pesticide concentration of 59.9 mg/kg body weight at which cent percent mortality occurred for 24 hours was taken as lethal dose value (24 hours, LD₁₀₀).

Calculation of LD₅₀ value

The LD₅₀ was estimated by the method of Litchfield and Wilcoxon (1949). The LD₅₀ value was obtained by the straight line interpolation method based on observed percentages of test animals surviving at concentrations lethal to more than half and less than half of the test animals. The LDC₀ value was derived by plotting the experimental data on a one scale log x probability sheet taking test concentration on the probability scale. A straight line was drawn between the points representing the mortality percentages Vs concentration (Standard methods published by American Public Health Association, 1960). From the point at which this line intersects 50 percent mortality, a perpendicular line was drawn to the concentration marked in ordinate and this indicate the LD₅₀ of 120 hours exposure period.

Estimation of blood glucose

Blood glucose was determined by the method of Murrel and Nace (1958). The Blood samples (0.01ml) were collected by caudal puncture using heparinized hypodermic syringe and were immediately deproteinized in 10 ml of 10 per cent tungstic acid. The solutions were filtered and the filtrate was used for glucose estimation. 0.5ml of the filtrate and 0.5ml of dilute tungstic acid were taken in a clean test tube and then 1.0ml of potassium ferricyanide solution was added. The test tube was placed in boiling water bath for 15 seconds and cooled in running tap water, when the contents were sufficiently cooled, 1.0ml of cyanide carbonate solution (buffer solution) was added and the tubes were again placed on boiling water bath for 15 minutes and then quickly cooled to 25-30°C. Then into each tube, 2.0ml of ferric dupanol reagent was added followed by the addition of 6.0ml of distilled water and the solution was mixed well and shaken well after 10 minutes the intensity of blue colour formed was read at 630 nm in a spectrophotometer against the reagent blank. A standard curve was constructed from the absorbance of the standard glucose solution. The unknown glucose values of blood samples were directly read from standard curve. The glucose values are expressed as mg/ml of blood.

Statistical analysis

The data obtained from the control and experimental parameters were analysed to determine the level of significance at various exposure periods by student 't' test.

3.RESULTS.

Biochemical studies

The mean blood glucose values of the controls and the median lethal dose treated lizards are presented in the Table 3 and illustrated in the Table 1. The median lethal dose treated lizards had decreasing trend in their blood of lizards, after administering with the pesticide, at different hours endosulfan viz., 6, 12, 18, 24, 48, 72, 96 and 120 were 103.25 ± 7.53 mg/100 ml, 99.95 ± 8.51 mg/100 ml, 96.86 ± 7.41 mg/100 ml, 93.26 ± 4.60 mg/100 ml, 92.19 ± 5.91 mg/100 ml, 91.22 ± 5.91 mg/100 ml, 79.89 ± 6.54 mg/100 ml and 78.17 ± 4.81 mg/100 ml of blood, while the control values were 115.27 ± 5.63 mg/100 ml, 114.40 ± 5.26 mg/100 ml, 114.83 ± 5.73 mg/100 ml, 114.87 ± 6.75 mg/100 ml, 115.10 ± 6.51 mg/100 ml, 115.50 ± 7.35 mg/100 ml, 114.50 ± 6.92 mg/100 ml and 114.64 ± 5.45 mg/100 ml of blood during the respective hours. The difference between the control and the treated lizards were appeared to be significant during all the experimental hours and the 't' value of different hours viz., 6, 12, 18, 24, 48, 96, and 120 as 3.39, 2.76, 3.29, 4.70, 5.89, 5.83, 8.65 and 11.94 ($P < 0.05$) respectively

Table: 1 Level of blood glucose (mg/100ml) of Chameleon zelenicus in the control and exposed animals Endosulfan at median lethal dose

Hours of administration	Control	Endosulfan exposure
6	115.28±5.63	103.25±7.53*
12	114.45±6.26	99.95±8.51*
18	114.83±6.73	96.84±7.41*
24	114.88±7.75	93.26±4.60*
48	115.10±7.51	92.19±5.91*
72	115.50±7.35	91.22±5.91*
96	114.50±6.92	79.89±6.54*
120	114.65±5.45	78.17±4.81*

Mean+S.D of six individual observations

*Indicates statistically significant at 5% level

Table:-2 Level of blood glucose (mg/100ml) of Chameleon zelenicus in the Control and Endosulfan exposed

Hours of administration	Control	Endosulfan exposure
6	115.28±5.63	99.95±8.32*
12	114.40±6.26	93.71±8.36*
18	114.84±6.73	87.17±7.86*

Mean+S.D of six individual observations

*Indicates statistically significant at 5% level

The mean blood glucose values of the control and the lethal dose treated lizards are presented in Table 4 and illustrated in the Table 2. The lethal dose treated lizards had decreasing trend in their blood glucose levels when compared to the control values upto 18 hours and later they died. The blood glucose level of lizards, after administrating with the pesticide endosulfan, at different hours viz 6, 12 and 18 were 99.95+8.36 mg/100ml and 93.85 +7.86 ml/ 100ml of blood, while the control values were 115.28 ± 5.63 mg/100 ml, 114.40 ± 5.26 mg/100 ml and 114.83 ± 5.73 mg/100 ml of blood during the respective the hours. The difference between control and treated lizards were statistically significant during all the experimental hours and their 't' values were 5.09, 6.12 and 8.65 (P<0.05) at 6, 12, and 18 hours respectively.

4.DISCUSSION

Blood glucose has been a sensitive biochemical indicator of environmental stress persuaded by handling, forced activity, thermal shock, contact with pesticides and other chemical pollutants (Chavin and Kavacevic⁸, 1961; Nakano and Tomlinson³⁷, 1967; Hill and Fromm²⁰, 1968; Watson and Mckeown, 1976; Wedemayer and Yasutaka, 1977 and Meielev et al., 1983). The blood glucose level which shows fluctuations represent a dynamic balance between the rate at which it enters the blood from the liver and the rate at which it is being removed by the body tissues from the blood for utilization (Saskin, 1941). In the present investigation, the normal blood glucose level was estimated in the laboratory acclimated lizards and its mean value was 115.5 + 7.35 mg/100 ml of blood. Khanna and Kumar²⁸ (1975) have stated that the lizards generally maintain a much higher level of blood glucose than do other reptiles. The present findings gain support from the findings of Dessauer¹⁶ (1952) for Gekko gecko, Dessauer¹⁴(1953) for Anolis carolinensis, Miller and Wurster (1958) for Eumeces obsoletus, Dawson¹² (1960) for Eumeces obsoletus, Dawson and Poulson¹³ (1962) for Phrynosoma cornutum, Motelica and Constanta (1965) for Lacerta vlridis viridis, Sabnis and Rangnekar⁴⁴ (1967) for Varanus monitor, Calotes versicolor and Nabuya carinata, Khanna and Kumar²⁸ (1974) for Uromastix hardwlccki and Madhavan and Rangnekar³⁰ (1985) for Calotes versicolor.

In the present study, it was documented earlier that the lizards administered with median lethal and lethal doses of endosulfan exhibited gradual decrease in their blood glucose level, when compared to control level. Similar results were reported for different species. Paul Ravindran⁴⁰ (1989) also recorded a decreasing trend in the blood glucose level of R. tigrina. Balakrishnan³ (1989) has been reported in the rat, Mllardla meltada administered with lindane showed the decreased blood glucose level. A decline in the blood glucose level was also found in the bird, Lonchura malacca administered with (endosulfan Usha,⁴⁸ 1990). Neelananarayanan³⁸ (1990) observed a similar decreasing trend in the blood glucose level in Mllardia mel tada when they poisoned with zinc phosphide. The present findings also gains support from the works of Mazeaud^{31,32} et al., (1971) for different fishes and aquatic organisms. Jawale²⁴ (1985) has observed for the freshwater fish, Rasbora daniconius on exposure to lethal concentration of the pesticides like DDT, endosulfan, rogor and dimecron, the fish showed a sudden fall in its blood glucose level which resulted in its death. Similar results were observed for Anabas testudineus administered with lethal concentration of has been endosulfan reported by Amudha² (1986). Mazeaud^{31,32} et al. (1971) have stated that when an animal was exposed to pesticides, which gives heavy stress on the animal and such stressful condition elicits neuro-endocrine responses in the organisms, which in turn affect the carbohydrate metabolism.

The rapid fall in the blood glucose level found in the present study might be due to the severe stress which occurred on the animal during the initial phase of anoxia. Wilhelmi⁵³ (1948) was of the opinion that under fully developed shock, the blood glucose level falls and then there is striking rise in the ratio of lactate to pyruvate and that the combination of an increased rate of utilization of carbohydrate. Engle¹⁷ (1949) has also stated that stress regularly results in hypoglycemia and death in cases where the adrenal cortex fails to function. It may be concluded that hypoglycemia in treated animals may be either due to failure of the adrenal to secrete the adrenocortical secretion or due to the damage caused in the adrenal due to the administration of the pesticide. The fall in the blood glucose level much below its normal threshold value resulted in hypoglycemic convulsions which might have caused the death of the treated animals while hyperglycemia has been reported for *C. versicolor* when administered with lindane (Thiyagesan⁴⁶, 1981) and when administered with endosulfan (Rajes wari⁴², 1988). They have accounted hyperglycemia as due to the increased secretion of glucocorticoids.

Blood glucose has been a sensitive biochemical indicator of environmental stress persuaded by handling, forced activity, thermal shock, contact with pesticides and other chemical pollutants (Chavin and Kavacevic,⁸ 1961; Nakano and Tomlinson³⁷, 1967; Hill and Fr



Chameleleo zelenicus (Females)

5. REFERENCES

1. **American Public Health Association. 1960.** Standard methods for the estimation of water and waste water. 11 edn. APHA, AWWA and WPCF, New York.
2. **Amudha, R. Christy. 1986.** Effects of malathion on blood glucose level and its histopathology on the endocrine adrenal gland of *Anabas testudineus* (Bloch). M. Phil., Thesis, Bharathidasan University
3. **Balakrishnan, G. 1989.** Effect of lindane (r - BHC) on the blood glucose metabolism and haematology of *Millardia meltada* (Gray). M.Phil., Thesis, Bharathidasan University.
4. **Banerjee, V. and M. Banerjee. 1969.** Seasonal variations of erythrocyte number with

- haemoglobin content in a common Indian lizard, *Yaranus monitor* (L). *Zool. Anz.*, 182 (3/4) : 203 - 207.
5. **Binyon, E.J. and G.I. Twigg. 1965.** Seasonal changes in the blood and thyroid of the grass snake, *Natrix natrix*. *Nature*, London. 207 : 779 - 780.
 6. **Calugrenan, M.L. 1970.** Effect of Cacl₂ on the cells of endocrine pancreas in the snake, *Natrix natrix*. *Horm Metab. Res.*, 2(2) : 94 - 101.
 7. **Chan, D.K.O., I.P. Callard and Chester Jones, I. 1970.** Observations on the water and electrolyte composition of the Iguanid lizard, *Dipsosaurus dorsal is* (Baird and Girard) with special reference to the control by the pituitary gland and the adrenal cortex. *Gen. Comp. Endocrinol.*, 15 : 374 - 387.
 8. **Chavin, W. and Kovacevic, A. 1961.** Adrenocortical histochemistry of intact and hypophysectomized gold fish. *Gen. Comp. Endocrinol.*, 1 : 264 - 274.
 9. **Choubey, B.J. 1975.** Seasonal changes in the blood of Indian lizard, *Calotes versicolor* (Daud,) *Zool. Anz.*, 194:: 35 - 41.
 10. **Conte, E.D. 1977.** Contiguity of the adrenaline-storing chromaffin cells with the interrenal tissue in the adrenal gland of a lizard. *Gen. Comp. Endocrinol.*, 32: 1 - 6.
 11. **Corbett, J.R. 1973.** The biochemical mode of action of pesticides. Academic Press, London, New York. pp : 107 - 165.
 12. **Dawson, W.R. 1960.** Physiological responses to temperature in the lizard, *Eumeces obsoletus*. *Physiol. Zool.*, 33: 87 - 103.
 13. **Dawson, W.R. and Poulson, T.L. 1962.** Oxygen capacity of lizard bloods. *Am. Midi. Nat.* 68 : 154 - 164.
 14. **Dessauer, H. C. 1953.** Hibernation of the lizard, *Anolis carolinensis*. *Proc. Soc. exp. Biol. Med.*, 82 : 351 - 353.
 15. **Dessauer, H. C. 1953.** Hibernation of the lizard, *Anolis carolinensis*. *Proc. Soc. exp. Biol. Med.*, 82 : 351 - 353.
 16. **Dessauer, H.C. 1952.** Biochemical studies on the lizard, *Anolis carolinensis*. *Proc. Soc. exp. Biol. Med.*, 80 : 742 - 744.
 17. **Engle, F.L. 1949.** *Endocrinology*. 45: 170 - 177.
 18. **Ghosh, A., G. Asok and B. Dutta. 1969.** Effect of reserpine on the release of adrenomedullary catecholamines and their role in glycemic and pressor responses in two avian species. In: "Progress in comparative endocrinology" (M.R.N. Prasad, Ed.). Suppl. 2 : 354 - 3 57.
 19. **Hague, R. and V.H. Freed. 1975.** "Environmental Dynamics of pesticides Plenum Press, New York and London.
 20. **Hill, C.W. and P.P. Fromm. 1968.** Response of the adrenal gland of rainbow trout, *Salmo galrdneri* to stress. *Gen. Comp. Endocrinol.*, 11 : 69 - 77.
 21. **Houssay, B.A. and J.C. Penhos. 1960.** Pancreatic diabetes and hypophysectomy in the snake, *Xenodon merremii*. *Acta Endocrinol.*, 35 : 313 - 323.
 22. **Hussaini, A. and A.H. Amani. 1976.** The effect of alloxan on the pancreatic islet tissue and blood sugar level of the toad, *Bufo regularis Reuss* and the lizard, *Agroa stellio*. *L. Zool. Soc. Egypt Bull.*, 28 : 11 - 12.
 23. **Hutton, K. E. and C.J. Goodnight. 1957.** variations in the blood chemistry of turtles under active and hibernating conditions. *Physiol. Zool.*, 30 : 198 - 207.
 24. **Jawale, M.D. 1985.** Effect of pesticide on metabolic rate of freshwater fish, *Rasbora daniconius*. *Environ. Ecol.*, 3 : 521 - 523.
 25. **Jayaraman, S. 1976.** Modulation of humoral and cell mediated immune responses to sheep erythrocyte in the lizard, *Calotes versicolor*. Ph.D., Thesis, Madurai Kamaraj University.
 26. **Kameswari, M. and L. Narsimha Rao. 1979.** Host-Parasite relationship : II. Post helminth infection haematological indices in *Calotes versicolor*. *Proc. Indian Acad. Sci.*, 88 ; 385 - 389.
 27. **Kekic, H. 1970.** A contribution to the knowledge of seasonal variation in some blood

- components of the lizard,
Lacerta viridis. *Jugol. Physiol. Pharmacol. Acta.*, 6(2) : 261 - 265.
28. **Khanna, S.S. and S. Kumar. 1975.** Blood glucose in the common Indian sand lizard, *Uromastix hardwicki*. *Copeia*, 4 : 767 - 771.
 29. **Lofts, B. and J.G. Phillips. 1965.** Some aspects of the structure of the adrenal gland in snakes. *J. Endocrinol.*, 33 : 327 - 328.
 30. **Madhavan, V. and p.v. Rangnekar. 1985.** Total pancreatectomy in the lizard, *Calotes versicolor*. *Indian J. exp. Biol.*, 23 : 523 - 524.
 31. **Mazeaud, D.P., J.A. Couch and A.M. Crane. 1971.** Insecticide susceptibility of some common fish family representatives. *Chesapeake Sci.*, 18 (1) : 677.
 32. **Mazeaud, D.P., J.A. Couch and A.M. Crane. 1971.** Insecticide susceptibility of some common fish family representatives. *Chesapeake Sci.*, 18 (1) : 677.
 33. **Mercier, M. 1981.** Criteria (Dose/Effect Relationships) for
 34. **Miller, M.R. and D.H. Wurstuer. 1958** Further studies on blood glucose and pancreatic islets of Lizards *Endocrinol.*, 63; 191-200.
 35. **Motelica, I. And M. Constant 1965 .** The action of insulin on the glycemia of certain reptiles, *Rev Rouma Biol Ser. Zool*, 9; 279-289
 36. **Murrel, L.R. and P.p. Nace. 1958.** Determination of glucose in fish blood. A modification of the Folin - Malmros micro-procedure. *Can. J. Biochem. Physiol.*, 36(11) : 1121 - 1124.
 37. **Nakano, T. and N. Tomlinson. 1967.** Catecholamines and carbohydrate concentrations in rainbow trout, *Salmo gairdneri* in relation to physical disturbance. *J. Fish. Res. Board Can.*, 24 : 1701 - 1715.
 38. **Neelanarayanan, P. 1990.** Laboratory studies of the effect of zinc phosphide on the biochemistry, histology and haematology in selected tissues of the soft furred rat, *Millardia melitana* (Gray). M.Phil., Thesis, Bharathidasan University.
 - 39 **Negilski, D.S. 1975.** Fundamental of aquatic bio assays. *Proc. R. Austr. Chem. Inst.*, 42 : 50 - 54.
 - 40 **Paul Ravindran, A.S. 1989.** Effects on malathion on the islets of Langerhans, adrenal glands and blood glucose level of the Indian Bull frog, *Rana tlgrina* (Daudin). M.Phil., Thesis, Bharathidasan University
 - 41 **Watson and M** level of the Indian Bull frog, *Rana tlgrina* (Daudin). M.Phil., Thesis, Bharathidasan University ckeown, level of the Indian Bull frog, *Rana tlgrina* (Daudin). M.Phil., Thesis, Bharathidasan University 1976;
 - 42 **Wede Rajeswari, A. 1988.** Toxicity, histopathology and physiological response in the female garden lizard, *Calotes versicolor* (Daud.) to malathion. M.Phil., Thesis, Annamalai University.
 - 43 **Rhoten, B. William. 1971.** Light and electron microscopic studies on panereatic islets of the lizard, *Lygosoma laterale*. *Gen. Comp. Endocrinol.*, 17 : 203 - 219.
 - 44 **Sabnis, P.B. and P.V. Rangnekar. 1967.** Some observations on the morphology of pancreatic islets and fasting blood sugar levels in lizards with particular reference to varanus monitor. *J. Biol. Sci.*, 10 : 14 - 19.
 - 45 **Silbergeld, E.R. 1974.** Blood glucose; A sensitive indicator of environmental stress in fish. *Bull. Environ. Oontam. Toxicol.*, 11 : 20 - 25.
 - 46 **Thiyagesan, K. 1981.** Pesticide (Lindane) as a stress factor in *Calotes versicolor*. M.Phil., Thesis, Annamalai University.
 - 47 **Unsicker, K. 1974.** Innervation of adrenal cells in the lizards, *Lacerta dugesi* and *Lacerta pityusensis*. *Gen. Comp. Endocrinol.*, 29 : 409 - 412.
 - 48 **Mayer and Yasutaka, 1977 and Usha, S. 1990.** Effects of endosulfan (35 EC) on the histological, biochemical and haematological aspects of the black headed munia (*Lonchura malacca*). M.Sc., Thesis, Bharathidasan University
 - 49 **Watson, T.A. and B.A. Mckeown. 1976.** The effect of sublethal concentration of zinc growth and plasma glucose levels in rainbow trout, *Salmo gairdnerii* (Richardson). *J.*

Wildlife Disease, 12 : 263 - 270.

- 50 **Watson, T.A. and B.A. Mckeown. 1976.** The effect of sublethal concentration of zinc growth and plasma glucose levels in rainbow trout, *Salmo gairdnerii* (Richardson). *J. Wildlife Disease*, 12 : 263 - 270.
- 51 **Wedemayer, G. 1972.** Some physiological consequence of handling stress in the juvenile Coho salmon (*Oncorhynchus kisutch*) and steel head trout (*Salmo gairdnerii*). *J.Fish. Res. Board Can.*, 29 : 1780 - 1783.
- 52 **Wedemayer, G.A. and W.T. Yasutaka. 1977.** Clinical methods for the assessment of the effects of environmental stress of fish health. Fish and wildlife Sci. Tech. Paper, 89, U.S.D.I., Washington.
- 53 **Wilhelmi, A.E. 1948.** Adrenal cortex. *Ann. Rev. Physiol.*, 10 : 259 - 276.

Organochlorine pesticides. Commission of the European communities. Pergamon

Press, Oxford. 381 pp.