

Neurohistopathological Effects of Gentamycin on Inferior colliculus of Adult Albino Rat

S M Dawar Husain

Abstract

Getamicin, an aminoglycosidic antibiotic, is ototoxic, nephrotoxic and causes neuromuscular blockade as well. A total of twenty albino rats (10 males and 10 females) were used in the present study, and they were equally divided into control and experimental groups. Experimental group rats received gentamicin intramuscularly for 21 days. Control group rats received normal saline. Then rats of both the groups were anaesthetized with nembutol, 35 mg/kg body wt. and perfused with 10% formalin. 10 μ thick sections of Inferior colliculus were stained with H&E and Glee's silver stains..

Observation under light microscope revealed degenerative changes.

Key words: Albino rats, Inferior colliculus, Gentamicin, Toxic effects

Introduction

Gentamicin is among the group of aminoglycosides that are used to treat aerobic gram negative bacterial infections. Amikacin, kanamycin, neomycin, streptomycin, paromomycin and tobramycin are other antibiotics in this group. The toxicity of these agents is dose related. Aminoglycosidic antibiotics block neuromuscular junction¹. Gentamicin was introduced in 1958 by Weinstein. It is nephrotoxic, neurotoxic and ototoxic and its side effects include ringing in ears, hearing loss, tinnitus, dizziness and anuria. Study was conducted on pharmacokinetics and dosage requirement of gentamicin in 1640 patients receiving treatment of gram-negative infections (daily dose ranged from 0.5 to 25.8 mg/kg² The effects of gentamicin were studied on 1327 patients, of which 31 patients (2.3%) had significant ototoxicity³ The average frequency of cochlear toxicity for gentamicin was reported to be 8.3% and exact incidence of vestibule-ototoxicity as about 3%⁴. Disequilibrium and ataxia were noted as main symptoms of vestibulotoxicity⁵. The chronic toxicity was related to aminoglycoside-phosphoinositol binding⁶. Evidence of neurotoxicity due to gentamicin and other aminoglycosides is available⁷.

A biochemical basis for the inherited susceptibility to aminoglycoside ototoxicity, has also been reported⁸. Greater sensitivity of the auditory cortex to aminoglycosidic antibiotics as compared to the periphery (cochlea) was reported⁹ Gentamycin toxicity was reported to depend on other factors like: dose and kidney function, other potentiating medications, genetic susceptibility and age¹⁰.

Though, the effects on auditory cortex have been reported along with ototoxic effects but the neurohistological effects of gentamycin on Inferior colliculus have less well been documented.

So, the present study is aimed to have further insight into the effects of gentamicin on the histology of the Inferior colliculus, which may explain central cause of ototoxicity.

Material and methods

20 adult albino rats, with equal number of males and females and weighing approximately 130 gms, were used in the present study. They were divided into control and experimental groups. Each group was comprised of 10 rats with equal male and female ratio. Experimental group rats were injected with gentamycin, 135mg/kg of body weight, intramuscularly for 21 days (Gentamycin WHO food Additives series 4, www.inchem.org/documents). Control group rats were treated with normal saline in same volume by intramuscular route for 21 days. After this duration, rats were anaesthetized by injecting nembutol, 35 mg/kg body wt and perfused with buffered 10% formalin. Inferior colliculus tissue samples were obtained from the brain. Tissue samples were processed for paraffin embedding. Then 10 μ thick sections were obtained with rotatory microtome. Sections were stained with H&E and Glee's silver stains for observation under light microscope.

Observations

In Haematoxylin & Eosin stained sections, control group rats, (Fig 1), showed clearly identifiable nuclei, while the experimental group rats (Fig 2). Showed scanty nuclei separated by clear zones.

In Glee's silver stained sections, compared to the control group rats (Fig 3), in the experimental group (Fig 4), no remarkable histological changes were observed.




Fig 1




Fig 2




Fig 3




Fig 4

Conclusion

Exposure of rat to gentamicin for three weeks produces no demonstrable microscopic changes in the Inferior colliculus..

References

1. C Pittenger, R Adamson. Antibiotic blockade of neuromuscular function. Annual review of pharmacology. 1972; 12, 169-184.
2. DE Zaske, RJ Cipolle, JC Rotschafer, DL Solem and RJ Strate. Gentamycin pharmacokinetics in 1640 patients: method for control of serum concentration. Antimicrobial agents and chemotherapy. 1982; 407-411.
3. G Arceiri, GG Jackson Ototoxicity of gentamycin in man: a survey and controlled analysis of clinical experience in the United States. J Inf Dis. 1971; 124 (suppl0, 130-137).
4. G Kahlmeter, JI Dahlager. Aminoglycoside toxicity: a review of clinical studies published between: 1975 to 1982. J Antimicrob Chemother. 1984; 13 Suppl A, 9-22.
5. A Takada and J Schacht. Calcium antagonist and reversibility of gentamycin induced loss of cochlear microphonics in the guinea pig. Hear Res. 1982; 8, 179-186.
6. MG Ganesan, ND Weiner, J Schacht. Effects of calcium and neomycin on phase behaviour of phospholipids bilayers. J pharma Sci. 1983; 72,1465-1466.
7. NA Faruqi, HS Khan. Effect of streptomycin and kanamycin on central nervous system: an Experimental Study. Indian Journal of Experimental Biology. 1986; 24, 97-99.
8. M Guan, N Fischel-Ghodsian, G Attardi. A biochemical basis for the inherited susceptibility to aminoglycoside ototoxicity. Human Mol Gen. 2000; 9, 12, 1787-93
9. VP Fissenko and NM Gusseinov. Electrophysiological study of the ototoxic effect of aminoglycoside antibiotics in freely moving animals. Vestin Otorhinolaryngol, Moscow Sachenov Medical Academy. 2003.
10. TC Hain. Gentamycin toxicity. Dizziness and Blance.com. 1999
11. AS Fix Ultrastructural pathology of the nervous system. V111 international symposium of the society of toxicologic pathologists meeting abstracts: “ toxicologic pathology of nervous system”. Toxicologic pathology of nervous system. Toxic pathol. 1999; 27, 690-704.
12. M Aschners, LG Costa. The reactive astrocytes. The role of Glia in Neurotoxicity. 2004; 2nd edition, 74.

Acknowledgement: We acknowledge the cooperation of technical and other supporting staff, Department of Anatomy, .J.N. Medical college, AMU Aligarh.